



**aesthetics**

Product Catalogue





## Mission Statement

Aesthetics Ltd is composed of a team of professionals with a mission to deliver innovative affordable and quality animal health products and agrochemicals to the COMESA region

## Our Vision

To be the leading source of quality and affordable animal health products and agrochemicals in the COMESA region.



*The Ministry of Environment and Natural Resources*

*in conjunction with the*

*United Nations Environment Programme-Ozone Secretariat*

*highly appreciates and recognises the efforts of*

*Aesthetics Limited*

*to meet the requirements of the Montreal Protocol to phase out*

*Ozone Depleting Substances in Kenya*



*Dr. Mohammed Isahakia*

*Parliamentary Secretary  
Ministry of Environment and Natural Resources  
Nairobi, Kenya*

*16 September 2000*

*Michael Gruber*

*Mr. Michael Gruber  
Deputy Executive Secretary  
Ozone Secretariat  
United Nations Environment Programme  
Nairobi, Kenya*

*16 September 2000*

# Products





# Super Skajcycline

## Contents

Oxytetracycline and Vitamins:

Nicotinic Acid	500mg	Vitamin D3	100,000i.u
Folic Acid	20mg	Vitamin E	200mg
Vitamin C	250mg	Vitamin K3	0mg
B.H.T	2.5mg	Vitamin B1	30mg
Copper	75mg	Vitamin B2	100mg
Oxytetracycline Hydrochloride	5.5g	Vitamin B6	65mg
Vitamin A	400,000i.u	Vitamin B12	3mg

Electrolytes include Potassium and Sodium Chloride and Acetate Salts

## Indication

1. In Disease
2. In Stress
3. To Day Old Chicks
4. More Eggs!
5. More Meat Gain!

Broad Spectrum antimicrobial. For:

1. Alimentary Tract Infections
2. Respiratory infections
3. Urinogenital infections
4. Septicaemias infections of unknown etiology where broad spectrum antimicrobial therapy is indicated.

## Usage instructions

Poultry of ALL ages:

One level teaspoonful per 5 litre drinking water for 5-7 days

Calves and Foals: Up to 50kg: 2-3 level teaspoonful's THREE times daily for a minimum of 3 days.

Over 50kg: 3-5 level teaspoons TWICE daily for a minimum of 3 days.

Lambs kids and Pigs.

Per 10kg: 2 level teaspoonful's TWICE daily for a minimum of 3 days.

100gm tub per 50kg of dry food for a minimum of 3 days.

## Pack size

100gm & 30gm

## Precautions

Withdrawal Period:

3 days for poultry Meat and Eggs.

7 days for other animal's meat and milk. Keep out of reach of children.

Store in a cool dry place





## Skavit

### Contents

Folic acid	200mg	Vitamin A	400,000 i.u
Calcium Panthothenate	300mg	Vitamin D	100,000 i.u
Vitamin C	250mg	Vitamin E	200mg
B.H.T	2.5mg	Vitamin K	30mg
Copper	75mg	Vitamin B	30mg
Cobalt	20mg	Vitamin B2	100mg
Zinc	550mg	Vitamin B6	65mg
Manganese	500mg	Vitamin B12	3mg
Iron	550mg	Nicotinic Acid	550mg

Electrolytes include: Potassium, Sodium chloride and Acetate salts

### Indication

#### Usage instructions

In drinking water:

Poultry: 1 level teaspoonful in 5 litres, daily for 5-7 days

Lambs, Kids and Pigs: 2 level teaspoonful's per 10kg bodyweight daily for 5-7 days

Calves and Foals: up to 50kg bodyweight- 2 level teaspoonful's, twice daily for 2 days.

Over 50kg- 1 level teaspoonful twice daily for 3 days

Cattle and Horses: 50g (half tub) daily for 3 days

In feed: 100g in 50kg. feed for one week.

#### Pack size

100g

#### Precautions

Store in a cool, dry place. Keep out of reach of children.



## SKAZONE®

### Contents

Sulphamethoxazole 10% w/w & Trimethoprim 2% w/w

### Indication

This is a water soluble Antibiotic.

It can be used for the control of infections caused by micro-organisms susceptible to Trimethoprim and Sulphamethoxazole such as bacterial and protozoal enteritis and urinary tract infections, Staphylococci infections in the ear and skin, salmonellosis and E. Coli infection in pigs, veal calves and poultry, coccidiosis, poultry typhoid, paratyphoid, giardiasis in poultry, respiratory tract infections and abscesses.

### Usage instructions

As directed by the Veterinarian.

General guidelines: To be administered twice daily in a little feed or water.

Poultry: One teaspoonful (5g) in 5 litres of drinking water for 5-7 days as a treatment dose. Half teaspoonful (2.5g) in 5 litres of drinking water for 14 days for prophylaxis. Calves, sheep, goats & pigs: One teaspoonful (5g) per 8kg body weight with water or feed for 5-7 days.

#### Pack size

100 g

#### Precautions

Water should be readily available to animals receiving this therapy. Do not use in animals hypersensitive to sulphonamides or those with liver disease.

Do not administer to pregnant animals.

Store in a dry place below 30°C. Keep out of reach of children.





## Oxycycline Pink Eye Treatment

### Contents

2.5% Oxytetracycline

### Indication

For the treatment of pink eye in cattle.

### Usage instructions

Apply once daily to the eyes for up to 7 days. Protect animals from direct sunlight, flies and dust.

### Pack size

50 g

### Precautions

Store in a cool, dry place. Keep out of reach of children.



## Epsom Salt

### Contents

Epsom salt

### Indication

As a sheath cleaner for breeding stallions.

Prevention of Grass tetany (in cows), and Grass staggers (in sheep) Relieves pain and inflammation for bruised or injured extremities. As a cathartic and laxative agent.

### Usage instructions

Dog: 2-25g

Cat: 2-5g

Horse: 30-100g

Cow: 250-500g

Pigs: 25-125g

Give orally with adequate drinking water during treatment.

### Pack size

500g & 250g

### Precautions

Store in a cool, dry place. Keep out of reach of children.





## Crak Milking Salve

### Contents

Dichlorophene 0.55% w/w in a Lanolised base

### Indication

For smooth and hygienic milking of cows.

Prevents and heals sores and cracks, of udders and teats, which are the predisposing factors of MASTITIS.

### Usage instructions

Apply to the hands as well as to the udders and teats of the cow, for a smooth and clean milking process.

### Pack size

400g, 250g, 100g

### Precautions

Store in a cool and dark place below 30°C . Keep out of reach of children.



## Healing Oil

### Contents

TetradecylTrimethyl Ammonium 0.5% w/v

### Indication

For the healing of abrasions, cuts, wounds and scratches of any animal..

### Usage instructions

Apply undiluted.


### Pack size

250ml & 100ml

### Precautions

Do not inject or drink.

Store in a cool, dry place away from sunlight. Keep out of reach of children



## Albanaex

### Contents

Albendazole 25mg/ml. 2.5%W/V

### Indication

For the treatment and prophylaxis of worm infections in cattle, calves, goats and sheep like: gastrointestinal worms, lungworms, tapeworms and liver flukes

### Usage instructions

Oral administration only. 120ml is enough to treat one adult cow or 10 adult sheep or goats.

Cattle: for every 10kg use 4ml

Calves: 40ml/100kg Young: 60ml/150kg Adult: 120ml/300kg

Sheep and Goats: Kids: 6ml/10kg Young: 8ml/20kg Adult: 10ml/30kg.

### Pack size

1 Litre, 500ml, 250ml & 120ml

### Precautions

Withdrawal period: Milk: 3 days

Meat: 7 days

Store in a cool dry place. Keep out of reach of children



## Piperex

### Contents

Piperazine Citrate BP, equivalent to 0.35g of Piperazine

### Indication

This is a water soluble WORM POWDER.

For the control of roundworms in poultry, pigs and calves.

Also for the for control of ascarids. Nodular worms. Pin worms and strongyles in horses.

### Usage instructions

Dissolve the entire contents of the 30g sachet in 20-30 litres of water per 100 birds. Or 1 ½ level teaspoonful's in 3-5 litres of water for every 15 birds.

Calves (100-150kg):

Dissolve the contents of one sachet in ½ litre of water and give as a drench.

Note: It is advisable to deworm at least every 3 months

### Pack size

10×30g sachets

### Precautions

Store in a dry place below 30°C. Keep out of reach of children



## Lasole

### Contents

Levamisole hydrochloride BP 15mg/ml

### Indication

Broad spectrum anthelmintic, used in the control and treatment of gastrointestinal worms and lungworm infections in cattle, calves and sheep.

### Usage instructions

5ml per 10kg body weight.

### Pack size

1 Litre & 125ml

### Precautions

Safe to use during pregnancy and lactation. Can also be used during vaccinations and other treatments simultaneously.

Store in a cool, dry place. Keep out of reach of children



## Albanaex+

### Contents

Albendazole 100mg/ml

### Indication

For the treatment and prevention of worm infections in cattle, calves, sheep and goat like: gastrointestinal worms, lungworms, tapeworms and liver flukes.

### Usage instructions

Round worms and tapeworms:

Sheep, goats, calves: 2.5ml for 50kg body weight. Cattle: 5ml for 50kg body weight.

Adult liver flukes:

Sheep, Goats, Calves: 5ml for 50kg body weight

Cattle: 10ml for 100kg body weight.

### Pack size

1 Litre, 500ml, 250ml, 120ml & 40ml

### Precautions

Withdrawal period:

Milk: 3 days

Store in a cool, dry place. Keep out of reach of children



## Aesbazole

### Contents

Albendazole 25mg/ml. 2.5%W/V

### Indication

For the treatment and prophylaxis of worm infections in cattle, calves, goats and sheep like: gastrointestinal worms, lungworms, tapeworms and liver flukes.

### Usage instructions

Oral administration only. 120ml is enough to treat one adult cow or 10 adult sheep or goats.

Cattle: for every 10kg use 4ml

Calves: 40ml/100kg

Young: 60ml/150kg

Adult: 120ml/300kg

Sheep and Goats:

Kids: 6ml/10kg

Young: 8ml/20kg

Adult: 10ml/30kg

### Pack size

1 Litre, 500ml, 250ml & 120ml

### Precautions

Withdrawal period: Milk: 3 days

Meat: 7 days

Store in a cool dry place. Keep out of reach of children.



## Aesbazole+

### Contents

Albendazole 100mg/ml

### Indication

For the treatment and prevention of worm infections in cattle, calves, sheep and goat like: gastrointestinal worms, lungworms, tapeworms and liver flukes.

### Usage instructions

Round worms and tapeworms:

Sheep, goats, calves: 2.5ml for 50kg body weight. Cattle: 5ml for 50kg body weight.

Adult liver flukes:

Sheep, Goats, Calves: 5ml for 50kg body weight

Cattle: 10ml for 100kg body weight.

### Pack size

1 Litre, 500ml, 250ml, 120ml & 40ml

### Precautions

Withdrawal period:

Milk: 3 days

Store in a cool, dry place. Keep out of reach of children.



## Albanaex 2500mg

### Contents

Albendazole 2500mg per bolus

### Indication

For the treatment and prevention of worm infections in cattle, calves, sheep and goat like: gastrointestinal worms, lungworms, tapeworms and liver flukes

### Usage instructions

Worm Dose: 7.5mg/kg Body weight

Fluke Dose: 10mg/kg Body weight

Body Weight	Number of Boli	
	Dose for worm	Dose for Fluke, Worm
100kg	¼	½
100 - 200kg	½	¾
200 - 250kg	¾	1
250 - 300kg	1	1¼
Over 300 Kg	1¼	1½

### Pack size

1 pack of 40 Boli

### Precautions

Do not exceed the stated dose Withdrawal period from treatment date: Milk: 3 days

Meat: 14 days

Store in a cool, dry place. Keep out of reach of children.



## Albanaex 600mg

### Contents

Albendazole 600mg per bolus

### Indication

For the treatment and prevention of worm infections in cattle, calves, sheep and goat like: gastrointestinal worms, lungworms, tapeworms and liver flukes.

### Usage instructions

Worm Dose: 7.5mg/kg Body weight

Fluke Dose: 10mg/kg Body weight

Body Weight	Number of Boli	
	Dose for worm	Dose for Fluke, Worm
Up to 50kg	½	1
100kg	1½	2
150kg	2	2½
200kg	2½	3½
250kg	3½	4½
300kg	4	5
350kg	4½	6

### Pack size

1 pack of 48 Boli

### Precautions

Do not exceed the stated dose.

Withdrawal period from treatment date: Milk: 3 days, Meat: 14 days

Store in a cool, dry place. Keep out of reach of children



## Albanaex 300mg

### Contents

Albendazole 300mg per bolus

### Indication

For the treatment and prevention of worm infections in cattle, calves, sheep and goat like: gastrointestinal worms, lungworms, tapeworms and liver flukes..

### Usage instructions

Worm Dose: 7.5mg/kg Body weight

Fluke Dose: 10mg/kg Body weight.

Body Weight	Number of Boli	
	Dose for worm	Dose for Fluke, Worm
Up to 50kg	1	1½
100kg	2½	3½
150kg	4	5
200kg	5	7
250kg	6½	8 ½
300kg	7½	10
350kg	9	11½

### Pack size

1 pack of 48 Boli

### Precautions

Do not exceed the stated dose

Withdrawal period from treatment date: Milk: 3 days, Meat: 14 days

Store in a cool, dry place. Keep out of reach of children



## Wormtrak 1 GM

### Contents

1 gram Levamisole Hydrochloride per bolus

### Indication

WORMTRAK 1 GM is a broad spectrum anthelmintic used in control and treatment of gastrointestinal worms and lungworm infections in cattle, calves, sheep and goats.

### Usage instructions

7.5mg Levamisole HCl per kg body weight or a bolus per 133 Kg body

Fluke Dose: 10mg/kg Body weight

Cattle /Calves		Sheep /Goats	
Body weight	Dose (Boli)	Body weight	Dose (Boli)
Upto 75 Kg	½	Upto 20 Kg	¼
100 – 150 Kg	¾ - 1¼	30 Kg	¼
200 – 250 Kg	1½ - 2	40 Kg	¼
250 – 300 Kg	2 - 2¼	50 Kg	¼
300 Kg & above	2½	60 Kg & above	½

### Pack size

1 pack of 32 Boli

### Precautions

Do not exceed the stated dose

Withdrawal period from treatment date: Milk: 3 days, Meat: 14 days

Store in a cool, dry place. Keep out of reach of children.



## Wormtrak 150mg

### Contents

Each Bolus Contains Levamisole HCL BP Vet 150 mg.

### Indication

WORMTRAK 150MG is a broad spectrum anthelmintic used in control and treatment of gastrointestinal worms and lungworm infections in cattle, calves, sheep and goats.

### Usage instructions

7.5mg Levamisole HCl per kg body weight or a bolus per 133 Kg body  
Fluke Dose: 10mg/kg Body weight.

Wt of Sheep, Goat & Pigs	Dosage (No of Tablets)
Upto 20 Kg	1
21-40 Kg	2
41-60 Kg	3
Above 60 Kg	5 (Maximum)

### Pack size

1 pack of 100 Boli

### Precautions

Do not exceed the stated dose.

Withdrawal period from treatment date: Milk: 3 days, Meat: 14 days

Store in a cool, dry place. Keep out of reach of children



## Oxycycline I.U Bolus

### Contents

Oxytetracycline Hydrochloride 2g per bolus

### Indication

For the treatment of bacterial infections after a difficult birth or retained placenta in cows, sows and ewes.

### Usage instructions

Cows: Insert one pessary only, two if the infection is severe

Sows and Ewes: Insert half a pessary only

### Pack size

10×2g boli

### Precautions

Store in a dry place below 25°C.

Protect from light

Keep out of reach of children



## Obgel

### Contents

Chlorhexidine gluconate and glycerin

### Indication

For rectal examination in Pregnancy diagnosis, obstetrical examination, artificial insemination, delivery and in day-to-day routine examination.

For lubrication of female genital tract during dystokia as well use to assist in normal delivery. In case of colic for removal of impacted dung. For digital examination of the rectum in animals which are small in size.

For lubrication of instruments at the time of genital examinations / treatment of genital infections.

For lubrication of thermometer bulbs at the time of recording temperature of new born foals / calves & small animals.

### Usage instructions

Liberal apply Obgel to adequately cover the area being lubricated.

### Pack size

1 Ltr x 12 btls

### Precautions

Store below 30°C protected from light in the original pack. keep out of reach of children



## Oxycycline Wound Spray

### Contents

Oxycycline Wound Spray

### Indication

- Foot rot
- Topical wound infection

### Usage instructions

Clean the area to be sprayed thoroughly

Pare (clip) hooves in foot rot

Spray the affected area from 6-8 inches away while holding the can upright.

Let the animal stand on dry ground for at least one hour

### Pack size

200 ml can x 12 x12

### Precautions

Avoid inhalation and contact with eyes

Avoid naked flames

Use in a well ventilated area.





## Biosafe

### Contents

15mg/ml Centrimide.

### Indication

A cationic detergent and antiseptic that penetrates all cracks and crevices, destroying bacteria and fungi.

### Usage instructions

Infectious disease outbreak: Three tablespoonful's in one gallon of water.  
Foot dip/ car wheel dip etc: Two tablespoonful's in one litre of water. General disinfection: One tablespoonful in one gallon of water. Wounds: Apply undiluted with cotton wool or cloth.

### Pack size

5 & 1 Litre

### Precautions

For external use only.  
Store in a cool, dry place.  
Keep out of reach of children



## Super Booster Foliar Fertiliser

### Contents

14% Nitrogen, 10% Phosphorous, 8% Potassium as well as trace elements and Growth hormones.

### Indication

A complete and balanced fertiliser that contains all the major macro elements, growth regulators and Amino Acids which are essential for optimal yields and healthy plant growth. It is specially formulated for RAPID ABSORPTION through the leaves.

### Usage instructions

Apply during the period of vegetative growth of the crop. It can be applied up to four times during the growth season.

### Pack size

1 Litre & 500ml

### Precautions

Wash hands and exposed skin after handling. Keep out of reach of children. Do not re-use the bottle. Do not spray into waterways or throw container into rivers.



## Super Blue Cross dust

### Contents

Fyfanon 2%w/w

### Indication

A Malathion based insecticidal dust for the CONTROL OF WEEVILS of maize beans and other grains.

### Usage instructions

50g/90kg of shelled maize, beans or other grains. Apply using a shovel. Sprinkle some dust over the packaging after filling to avoid reinfection. Repeat after 5 months.

### Pack size

200g, 400g, 1kg, 2kg, 5kg & 25kg

### Precautions

Do not apply the powder to surfaces where food is prepared or handled. Do not eat, drink or smoke while applying. Do not inhale the dust. Wear gloves and apron while applying, wash thoroughly after. Store in a cool, dry place. Keep out of the reach of children



## Vegetable dust

### Contents

Malathion (Fyfanon®) 3.5%w/w

### Indication

A Malathion based insecticidal dust that is effective against a wide range of insect pests on vegetables, such as: Aphids, Mites, Scales, Ballworms, and Leaf miners etc. On Cabbage, Kale, Celery, Tomatoes and other vegetable plants

### Usage instructions

Sprinkle the dust when pests are sighted on the vegetables, repeat after 10 days if necessary

### Pack size

400g

### Precautions

Do not apply the powder to surfaces where food is prepared or handled. Do not eat, drink or smoke while applying. Do not inhale the dust. Wear gloves and apron while applying, wash thoroughly after. Store in a cool, dry place. Keep out of the reach of children.



## New Pal Dog Shampoo

### Contents

D-Phenothrin (Sumithrin®) 0.4%w/w.

### Indication

A medicated shampoo for dogs that kills all ticks, fleas and lice leaving the coat silky and smooth.

### Usage instructions

Mix 100ml of the shampoo with 1 litre of water. Wet the dogs coat thoroughly with plain water then apply the shampoo mixture. Lather the dog's coat paying particular attention to the neck scruff, ears and between the toes. 100ml of shampoo is adequate for an adult and 50ml for puppies of three months or older..

### Pack size

5 Litres & 1 Litre & 250ml

### Precautions

Do not apply the liquid to surfaces where food is prepared or handled. Do not eat, drink or smoke while applying. Wear gloves and apron while applying, wash thoroughly after. Store in a cool, dry place. Keep out of the reach of children.



## Pal Pet Ticks & Flea Spray Aerosol

### Contents

D-Phenothrin (Sumithrin) 0.2% w/w.

### Indication

This is a Sumithrin based insecticidal spray that can eliminate external parasites like fleas, ticks and lice from dogs and cats.

### Usage instructions

It is effective against dog's external parasites such as fleas and ticks. Strong Killing Effect, Low Mammalian Toxicity, Good Stability and free from offensive odor.

Spray the PAL PET TICKS & FLEA SPRAY AEROSOL on the ear, scruff of the Neck, between the toes and behind the tail. Do not exceed spraying for more than 10 seconds per animal, which should discharge 10 ml of the liquid. Repeat once a week or as directed by your veterinary doctor

### Pack size

200 ml x 12 pcs

### Precautions

Do not use on Cats. Not recommended for puppies below Eight weeks. Store under lock and key in original, tightly closed containers in a cool , dry place away from food, feeds, children and domestic animals. Do not spray directly into the eyes of the dog.



## Pal Pet Powder

### Contents

Malathion (Fyfanon\*) 1% w/w

### Indication

This is a Malathion based insecticidal dust that can eliminate external parasites like fleas, ticks and lice from dogs and cats.

### Usage instructions

Dust the powder directly from the container by dusting coats of the animals, pay particular attention to sites of predation of ticks such as interdital spaces beneath tail, around the ears and scruff of the neck. Kennels and cat baskets and other sites which pets often inhabit should be treated.

Use 50gm per adult dog/cat. 10 gm for puppy/kitten greater than three months of age.

### Pack size

300 g

### Precautions

Do not apply powder on surfaces where food is prepared or handled.

Store under lock and key in original, tightly closed containers.

Store in a dry place below 30°C.

Keep out of reach of children.



## Poultry Dust

### Contents

Malathion (Fyfanon\*) 4% w/w

### Indication

It is effective against poultry external parasites such as fleas and poultry louse

### Usage instructions

Hold the bird by the legs turned upside down. Dust the powder directly from container to areas around the eyes, under the wings and the legs. Repeat when infestation is noted again. 50 gms is enough for one bird and 25gms for one chick aged two weeks or older.


### Pack size

200g x 24 pcs

400 g x 24 pcs

### Precautions

Store under lock and key in original, tightly closed containers in a cool, dry place away from food, feeds, children and domestic animals. Do not sprinkle directly into the eyes of the bird.



## Stop Vermin Powder

### Contents

Cyphenothrin (Gokilaht) 0.5% w/w.

### Indication

An insecticidal dust effective against household crawling insect pests such as cockroaches and ants.

### Usage instructions

200g/100 square feet of surface area. Dust the powder directly from the container onto the areas infested with vermin, including the cracks and crevices.

### Pack size

200g & 100g

### Precautions

This product is poisonous. Cover all utensils while dusting. Do not apply the powder to surfaces where food is prepared or handled. Do not eat, drink or smoke while applying. Do not inhale the dust. Wear gloves and apron while applying, wash thoroughly after. Store in a cool, dry place. Keep out of the reach of children and domestic animals



## Stalk Borer Treatment Powder

### Contents

Malathion (Fyfanon\*) 3.5% w/w.

### Indication

This is a Malathion based insecticidal dust effective against stalk borers in maize.

### Usage instructions

Sprinkle the dust into the growing point of the plant. Sprinkle the dust when the maize has between 6-7 leaves. One shake of container for one plant. Repeat fortnightly until ear formation.

### Pack size

400g x 12 pcs

### Precautions

Store under lock and key in original, tightly closed containers in a cool, dry place away from food, feeds, children and domestic animals. Do not sprinkle directly into the eyes of the bird.

# STOP!

## CRAWLING INSECTS



*Put irritating pests in their place ...*



From



**Stop vermin Powder** is a Cyphenothrin based insecticidal dust effective against household crawling insect pests such as cockroaches and ants.

**Available in 100gms and 200 gms**

P.O. Box 18171 - 00500, Nairobi. Tel: 6537615 / 6, 555157 Safaricom: 07 11 11 8148 Airtel: 0735 547 014  
email: [aesthetics@iconnect.co.ke](mailto:aesthetics@iconnect.co.ke) | [aestheticslimited@gmail.com](mailto:aestheticslimited@gmail.com) FAX: 552930

Veterinary  
Equipment  
&  
Surgical  
Instrument



# Dehorning

---



[Dehorning Wire 12 meters](#)

---



[Dehorning Wire Saw Handle](#)

---



[Dehorning Iron](#)

---



[Electric Dehorning Iron](#)

---



# Castration

Burdizo Castrator for Sheep - Small



Burdizo Castrator - Medium



Burdizo Castrator - Large



Plastic - Rubber Castration Ring Applicator



Castration Bands Orange 100s & 500s



Metallic - Rubber Castration Ring Applicator



Castration Knife



# Dairy



Milk strainer  
stainless steel 11 Ltr



Milk Strainer



Strainer inset perforated disc



Strainer inset meshing  
Ø170mm



Strip Cup



Latex Teat



Large Floating Teat



Milk Test Liquid  
KERBA TEST 1 Ltr



Dosing bottle 250 ml  
for milk test



Teat Dip



Milk Filter



Mastitis Indicator Paper




Lactometer



1&2 Ltr Calf Feeding Round/Square  
Bottle with latex Teat



Calf anti-sucker - Plastic  
with spikes



Calf Anti-sucker -  
Aluminium with spikes



Obstetrical Gloves



Milk Catheter



Milk Splitter



Hobble Chain for  
Milking

## Milk Cans



5 Ltr



10 Ltr



15 Ltr



20 Ltr



25 Ltr



30 Ltr



40 Ltr



50 Ltr

## Milk Buckets



5 Ltr




10 Ltr



15 Ltr

## Milk Scoops



1/2 Ltr



1 Ltr

## Measuring cup 1/2, 1 & 2 ltr Plastic



# Dairy

---



One Hand Hair Clipper



Sheep Shear



Stomach tube for Calves



Weight Measure



Thermometer



Electric Prodder KAWE



Stomach tube for Cattle

# Syringes & Accessories




Automatic Poultry  
Vaccinator



Hauptner Syringe



Roux Revolver Syringe



Roux Revolver & Hauptner Syringe  
Glass Barrel & Washers



## Hypodermic Needles

- Re-Usable Hypodermic Needles 14G X 1/2
- Re-Usable Hypodermic Needles 14G X 1
- Re-Usable Hypodermic Needles 16G X 1/2
- Re-Usable Hypodermic Needles 16G X 3/4
- Re-Usable Hypodermic Needles 16G X 1
- Re-Usable Hypodermic Needles 16G X 1 1/2
- Re-Usable Hypodermic Needles 17G X 1/2
- Re-Usable Hypodermic Needles 17G X 3/4
- Re-Usable Hypodermic Needles 17G X 1
- Re-Usable Hypodermic Needles 17G X 1 1/2
- Re-Usable Hypodermic Needles 18G X 1/2
- Re-Usable Hypodermic Needles 18G X 3/4
- Re-Usable Hypodermic Needles 18G X 1
- Re-Usable Hypodermic Needles 18G X 1 1/2

# Syringes & Accessories



Re-Usable Metallic Syringe - 10ml, 20ml, 30ml & 50ml



Re-Usable Metallic Syringe - Plastic, Steel & Copper Tip Injectors



Suture needles, set 1+4+7



Re-Usable Metallic Syringe - 100 ml Syringe

# Surgical Equipment



Thinning Scissors  
with Teething on both sides



Hair Scissors  
(with Rounded Tips)



Curved Scissors




Towel Forceps



Scissors Straight



Scissors Curved



Tooth Cutting Forceps



Mathieu Needle Holder



Tissue Forceps



Dissecting Forceps



Splinter Forceps

# Surgical Equipment



B.P Scalpel Blade




B.P Scalpel Blade Holder No.4



Ruminal Magnet



Post Mortem Tray



Birth Cord for Calf



Ceramic Ruminal Magnet



Tooth Nipper



Trocar & Cannula



Sewing material Perlon, size 3 thread



Post Mortem Kit




Sewing material size 3



Sewing material size 4



# Drenching



Automatic Drenching Gun  
available with Knapsack  
container and in manual models



Drenching Cannula



Balling gun



Automatic Model - 200ml / 60ml / 30ml



Manual Model I




Manual Model II

# Animal Identification


---



Ear Tags for cows, sheep and goats




Ear Tag for Rabbits



Ear Tag for Rabbits

# Animal Identification



Flexa Ear Tag  
Applicator

## Ear Tag Identification




Marking Pen for  
The Unmarked Tag



Prima Flex (Universal)  
Eartag Applicator


## Tattoo Identification



Tattooing Numbers  
and Letters



Tattooing Ink



Tattooing Pliers



Set of paint brands  
0-9, 10 cm



Tattoo ink roll-on,  
black




Horizontal Tattooing  
Forceps

# Animal Identification



Ear Notcher



Ear Notcher



Marking Sticks  
(Assorted Colours)



Livestock Marking  
Spray



Cold Branding Iron  
(0-9)



Master Laine Ink  
(for Cold Branding Iron)



Harness for Bulls and  
Rams



Marking Black for  
Ram & Bull Harness



Cow halter with chin  
chain



Sisal-PP halter  
without clamp

# Hoof & Claw Treatment



Hoof Knife



Hoof Claw Cutter  
Burdizzo Type



Foot Rot Hoof Shear



Foot Rot Hoof Shear  
with Pocket



Nail Claw Cutter



Sheep shear, 31 cm,  
double bow

# Restraining Equipment


---



Bull Nose Ring Stainless Steel  
Large size 3½ inch (88.9mm)  
Medium size 3 inch (76.2mm)  
Small size 2½ inch (63.5mm)



Bull Nose Ring Copper  
Large size 3½ inch (88.9mm)  
Medium size 3 inch (76.2mm)  
Small size 2½ inch (63.5mm)



Harm's Bull Holder

# Diagnostic Equipment



Single Line

Double Line Stethoscope



Double Line

## Thermometers



Clinical Thermometer



Digital Thermometer

# Pumps

---



Eva 2 Litre Pump



Alta Spray 10 Litre



Alta Spray 15 Litre



Conico Spray 1 Litre



# Externals





## Liquid Paraffin

### Contents

Contains liquid paraffin B.P.

### Uses

1. As a laxative in small quantities.
2. Eases irritation in eczema and nappy rash.
3. As an emollient lotion to treat/soothe dry skin.
4. As an ingredient in eye lubricants.

### Usage instructions

Apply generously on the skin. For laxative qualities, use 1 teaspoonful for infants and 1 tablespoonful for adults taken at night..

### Pack size

50 ml x 100 pcs per carton

100 ml x 100 pcs per carton

## Hydrogen Peroxide

### Contents

Contains hydrogen peroxide B.P 6% w/v (20 volume)..

### Uses

1. To disinfect cuts and stop bleeding on superficial cuts and grazes.
2. Treating/ removing flesh blood stains on clothing and other items.
3. Removal of organic odors (in mouth and feet and small dump sites) by oxidizing hydrogen sulphide.
4. Cleaning/ removal of grout from floors.
5. Provide oxygen to fish in small aquariums.

### Usage instructions

First aid: Dilute with equal volume of water.

To arrest bleeding and disinfect wounds apply diluted solution to lint or cotton wool and cleanse the wound.

Removal of wound dressing: Dilute with equal volume of water.

Soak dressing with diluted solution, leave for a few minutes, and remove the dressing. Mouth wash & Deodorant:

Dilute two teaspoonfuls to one glass of clean water and gargle.

### Safety Precautions

Irritant solution. Avoid contact with skin and eyes. In case of accidental splashing in eyes or skin, wash with plenty of water and seek medical advice.

Not to be taken by mouth unless diluted as indicated above.

### Pack size

200 ml x 25 pcs per carton

5 Ltrs x 4 pcs per carton





## Pure Glycerine

### Contents

Contains pure glycerine B.P.

### Uses

1. Glycerin is a humectant, (attracts moisture to the skin). This way it softens the skin on application since it is also an excellent emollient.
2. It is used as a base for skin care lotions.
3. Used as a stool softener when applied rectally.

### Usage instructions

Apply generously to the skin as desired. For application as a stool softener, introduce 5 ml through the rectum in the prone position.

### Safety Precautions

This product is for use on the outside of the body only. Do not apply over deep or puncture wounds, infections or lacerations unless directed by a doctor. Do not use on animal bite, serious burns. When. Do not get into eyes.

### Pack size

- 50 ml x 100 pcs per carton
- 100 ml x 100 pcs per carton



## Tincture of Iodine

### Contents

Contains 2.5% w/v Iodine

### Uses

1. General and disinfection
2. Disinfection of wounds & burns

### Usage instructions

Before use, clean the affected area, rinse well, and pat dry. Using a cotton swab to avoid staining your hands, apply gentian violet to the affected area, usually once or twice a day or as directed by the physician. Apply only to affected area and avoid surrounding healthy skin.

Wash hands immediately after use unless the area being treated includes hands. Avoid getting the product in your eyes. If medication gets into your eyes, rinse the area immediately with clean water for at least 15 minutes. Seek medical advice if your eye becomes irritated. Do not cover the area with plastic or waterproof bandages. Covering the area may increase the risk of skin irritation

### Safety Precautions

This product is for external use only. Do not apply inside the mouth, nose or ears unless directed by the physician. If the physician has prescribed this medication for area in the mouth, you/your child must be careful to avoid swallowing the medication.

This medication is a dye and will stain skin and clothing. Do not apply to open sores (skin ulcers). This may cause permanent 'tattooing' of the skin

### Pack size

- 50 ml x 100 pcs per carton
- 100 ml x 100 pcs per carton



## Gentain Violet

### Contents

Contains Gentian Violet 1% W/V

### Uses

1. Marking the skin for surgery preparation and allergy testing.
2. Tinea; e.g. Athlete's foot, jock itch, and ringworm.
3. Candida albicans and related infections; e.g. thrush, yeast infections, Mouth ulcers.
4. Abrasion and weeping wounds where the flesh is abraded away.
5. Prevents infection in almost any wound or burn and quickly help the epithelium form a scab.
6. As a tissue stain in the preparation of light microscopy sections.

### Usage instructions

Before use, clean the affected area, rinse well, and pat dry. Using a cotton swab to avoid staining your hands, apply gentian violet to the affected area, usually once or twice a day or as directed by your doctor. Apply only to the affected area, and avoid surrounding healthy skin.

Wash hands immediately after use unless the area being treated includes the hands. Avoid getting the product in your eyes. If the medication gets into your eyes, rinse the area immediately with clean water for at least 15 minutes. Seek medical attention if your eye becomes irritated.

Do not cover the area with plastic or waterproof bandages. Covering the area may increase the risk of skin irritation.

### Safety Precautions

This product is for use on the outside of the body only. Do not apply inside the mouth, nose, or ears unless directed by your doctor. If your doctor has prescribed this medication for areas in the mouth, you/your child must be careful to avoid swallowing the medication.

This medication is a dye and will stain skin and clothing. Do not apply to open sores (skin ulcers).

This may cause permanent "tattooing" of the skin.

### Pack size

50 ml x 100 pcs per carton

100 ml x 100 pcs per carton

## Surgical Spirit

### Contents

Surgical Spirit contains denatured alcohol 70% v/v

### Uses

1. Skin disinfection before administering an injection, after minor injury or before a surgical procedure.
2. Hand disinfection in sterile zones.
3. Hardening of sensitive skin especially in convalescent state such as bed sores in bedridden persons.
4. Cleaning and disinfection of surfaces and equipment in hospital set-ups.
5. Shaving equipment and Skin disinfection after shaving.

### Usage instructions

Use undiluted on intact skin, surfaces or instruments.

### Safety Precautions

Avoid contact with mucous membranes and damaged skin surfaces.

### Pack size

- 50 ml x 100 pcs per carton
- 100 ml x 100 pcs per carton
- 500 ml x 30 pcs per carton
- 1 Ltr x 30 pcs per carton
- 5 Ltr x 4 pcs per carton



## Methylated Spirit

### Contents

Methylated spirit contains denatured alcohol 70% v/v

### Uses

1. Skin disinfection before administering an injection or after a minor injury.
2. Shaving equipment and Skin disinfection after shaving.
3. As a fuel in light warming of food and lighting of pressure lamps.

### Usage instructions

Use undiluted on intact skin, surfaces, instruments or as a fuel.

### Safety Precautions

Avoid contact with mucous membranes and damaged skin surfaces.

### Pack size

- 50 ml x 100 pcs per carton
- 100 ml x 100 pcs per carton
- 500 ml x 30 pcs per carton
- 1 Ltr x 30 pcs per carton
- 5 Ltr x 4 pcs per carton





## Castor Oil

### Contents

Contains Castor oil BP

### Uses

1. Used as a stimulant and lubricating laxative.
2. Used for skin conditioning, such as in shampoo, lipstick and lip balm.
3. Used as a skin anti-inflammatory agent and antimicrobial.
4. Application to the lower abdomen, legs and hands, and in palms to reduce body heat.
5. Other uses include a variety of skin conditions like keratosis, dermatosis, wound healing, acne, ringworm, warts and other skin infections, sebaceous cysts, and itching. Castor oil also enhances the absorption of other agents across the skin.

### Usage instructions

Use undiluted by applying on the desired action points. For laxative effects, take 1 teaspoon twice daily and discontinue the therapy when the desired effect is achieved.

### Safety Precautions

Avoid contact with mucous membranes and eyes.

### Pack size

50 ml x 100 pcs per carton

100 ml x 100 pcs per carton



## Aeslon

### Contents

Contains Cetrimide 3% w/w, Chlorhexidine digluconate 0.3% w/v

### Uses

1. General and skin disinfection
2. Cleaning & disinfection of equipment
3. Disinfection of wounds & burns
4. Pre-operative disinfection of hands
5. Prevention of microbial colonization of peripheral intravenous catheters.

### Usage instructions

Dilute 60-90ml in 1-1.5 litres of water. Higher concentrations are indicated for use on surfaces.

### Safety Precautions

Avoid contact with mucous membranes and eyes

### Pack size

5Ltr x 4 pcs per carton





aesthetics  
animal health



aesthetics  
agro-chem



aesthetics  
poultry



aesthetics  
pet care

## **Aesthetics Ltd.**

P.O. Box 18171 - 00500, Nairobi, Kenya

House of Aesthetics, Pembe Road off Lusaka Road,  
Industrial Area

Tel: +254 20 555 157 / 653 7616 | Fax: +254 20 552 930

Email: [aesthetics@iconnect.co.ke](mailto:aesthetics@iconnect.co.ke)